

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.



March 2002

Prof. Uri Sivan
Department of Physics and Solid State Institute
Bertoldo Badier Academic Chair
Technion -- Israel Institute of Technology
Haifa, 32000 Israel
Tel: +972-4-8293452, Fax +972-4-8235107
Email: phsivan@tx.technion.ac.il

A. PERSONAL

Born: Haifa, Israel, 1955.
Domestic Status: Married + 3

B. EDUCATION

1979 - 1982 - B.Sc. in Physics and Mathematics - Tel Aviv University, *magnam cum laude*
1982 - 1984 - M.Sc. in Physics - Tel Aviv University, *magnam cum laude*
1984 - 1988 - Ph.D. in Physics, Tel Aviv University, *summa cum laude*

C. PROFESSIONAL EXPERIENCE

1988-1991 - IBM, T. J. Watson Research Center, NY, USA
1991-1995 - Senior Lecturer, Physics Dep. Technion-IIT, Israel
1995-1999 - Associate Prof., Physics Dep. Technion-IIT, Israel
1999-present - Professor, Physics dep., Technion - IIT, Israel.

D. INVITED TALKS IN INTERNATIONAL CONFERENCES-RECENT 5 YEARS

1995 Franco-Israeli meeting, Paris - France, *Mesoscopic Fluctuations in the Ground State Energy of a Disordered Quantum dot.*
1995 Aronov Memorial Symposium, Zichron Ya'akov - Israel, *Spectroscopy of Disordered Quantum Dots.*
1996 Conference on Quantum Chaos in Mesoscopic Systems, Santa Barbara - USA, *The chemical Potential of A Strongly Interacting 2D Fermion Layer.*

- 1996 Spring College in Condensed Matter Physics, International Center for Theoretical Physics, Trieste -Italy, three lectures course on *Experiments in Mesoscopic Systems*.
- 1996 Adriatico Research Conference on Mesoscopic Phenomena in Complex Quantum Systems, Trieste - Italy, *Excitation Spectrum and Ground State Level Statistics of Disordered Quantum Dots*.
- 1996 International Conference on "Correlated Electrons in Systems of Reduced Dimensionality", Trieste-Italy, *The Thermodynamics of Strongly Correlated Fermions*.
- 1996 International Conference on Electron Localization and Quantum Transport in Solids, Jaszowiec - Poland, *Experiments on Level Statistics in Diffusive Quantum Dots*.
- 1996 International Conference on Electron Localization and Quantum Transport in Solids, Jaszowiec - Poland, *The Compressibility of Fermions at High F_0 Numbers*.
- 1997 Minerva Conference on Mesoscopic Physics, Eilat - Israel, *The Coulomb Blockade Revisited*.
- 1997 Japan-Israel binational conference on mesoscopic physics, Beer Sheva - Israel, *The Role of Coulomb Correlations in Quantum Dots*.
- 1997 The annual meeting of the Israeli Physical Society, Beer Sheva - Israel, *Correlation Energy of Strongly Interacting Fermions at High F_0 Numbers*.
- 1997 International workshop on "Direction in Mesoscopic Physics", Leiden-Holland, *The Coulomb Blockade, Revisited*.
- 1997 International Conference on Mesoscopic Physics, MESO97, Chernogolovka - Russia, *Correlation Energy of Strongly Interacting Fermions at High F_0 Numbers*.
- 1997 International conference on "Fundamental Aspects of Applications of Single Electron Devices", Lyngby - Denmark, *Ground State Properties of a Quantum Dot*.
- 1997 International Conference on Strongly Coupled Coulomb Systems, Chestnut Hill - USA, *2D Fermions - Experiment* (presented by Dr. S. Shapira).
- 1998 International Summer School on Coherence in Electronic Systems, Ustron, Poland, *Coherence and Dephasing in Low Dimensional Systems*.
- 1998 European Conference on Mesoscopic Physics, PHASDOM-98, Neuchatel, Switzerland, *Self Assembly of Nanoscale Electronics by Biotechnology*.

-
- 1998 Nordic-Baltic Physical Society meeting, Nyborg, Denmark, *Self Assembly of Nanoscale Electronics by Biotechnology.*
 - 1998 International Conference on the Physics of Semiconductors (ICPS Jerusalem, Israel, *Coulomb Drag in the Quantum Hall Effect Regime.*
 - 1998 International Workshop on "Disorder and Interactions in Quantum Hall and Mesoscopic Systems", Santa Barbara, USA, *Coulomb Correlations in a Dilute 2D Fermion System.*
 - 1998 International Workshop on "Electron Transmission through Molecules and Interfaces", Zanzach, Israel, *Self assembly of Nanometer Scale Electronics using Biotechnology.*
 - 1999 The annual meeting of on the American Material Research Society (MRS), San Francisco USA, *Self Assembly of Nanometer Scale Electronics by Biotechnology.*
 - 1999 The International Conference on the Physics of Two Dimensional Electronic Systems-EP2DS 22, Ottawa Canada, *Self Assembly of Nanometer Scale Electronics by Biotechnology.*
 - 1999 International workshop on "New Developments in Quantum Hall Effect", Minneapolis USA, *Electron-Hole Drag in the QHE Regime.*
 - 1999 The German Physical Society Meeting, Munster, Germany, *Self Assembly of Molecular Scale Electronics by Biotechnology.*
 - 1999 Nato Advanced Research Workshop, Kiev, Ukraine, *Some Aspects of Molecular Self Assembly.*
 - 1999 Rencontres de Moriond, Les Arcs France, *Metal-Insulator Transition in 2D?*
 - 1999 The First Stig Lundqvist Research Conference on the Advancing Frontiers in Condensed Matter Physics - Quantum Phases in Electron Systems of Low Dimensions, Trieste Italy, *The Compressibility of Strongly Correlated 2D Fermions Near the Metal Insulator Transition.*
 - 1999 The "Microelectronics Advanced Research Initiative" workshop on Nanofabrication", Marseille France, *Self Assembly of Molecular Electronics by Biotechnology.*
 - 1999 IST99 Conference on "wet frontiers in microelectronics - the interface between biology and microelectronics", Helsinki Finland, *Conceptual and Practical Challenges in the Self Assembly of Microelectronics Using Biotechnology.*
 - 2000 Nato Advanced Research Workshop on "Frontiers in Nano-Optoelectronics Systems", Kyev Ukraine, *Self Assembly of Molecular Scale Electronics by Biotechnology.*

- 20/10 Winter School on the physics of low dimensional systems, Monteford Austria, *Molecular Electronics by Biotechnology*.
- 20/10 International workshop on "Chaos and Interaction in Quantum Dots", Minnesota, USA, *Condensation of Positively Charged Colloids on DNA*.
- 20/10 MesoSpin 2000, Cortina, Italy, *Condensation of Positively Charged Colloids on DNA*.
- 20/10 Japan-Israel binational meeting, Tokyo Japan, *Condensation of Positively Charged Colloids by DNA*.
- 20/11 10th Brazilian Workshop on Semiconductor Physics, Guarujá SP Brazil, *Self Assembly of Molecular Scale Electronics by Biotechnology*.
- 20/11 Rencontre de Moriond meeting, Les Arcs France, *Evidence Against Metal-Insulator Transition in Two Dimensional Holes*.
- 20/11 American Physical Society meeting, Seattle Washington USA, *Evidence Against Metal-Insulator Transition in Two Dimensional Holes*.
- 20/11 NATO summer school, Windsor England, *Self Assembly of Molecular Scale Electronics by Biotechnology*.
- 20/11 Workshop on Nanoscience, Dresden, Germany, *Summary and Perspectives of Nanotechnology*.

B. LIST OF PUBLICATIONS

1. J. Salzman, U. Sivan, E. Kapon, and A. Katzir, "Heterodyne Detection Using Multimode Waveguide Y-Couplers", *Appl. Opt.* **22**, 31 (1983).
2. A. Yariv, C. Lindsey, and U. Sivan, "Approximate Analytic Solution for Electronic Wavefunctions and Energies in Coupled Quantum Wells", *J. Appl. Phys.* **58**, 3669-3672 (1985).
3. U. Sivan and Y. Imry, "Multichannel Landauer Formula for Thermoelectric Transport with Application to the Thermopower Near the Mobility Edge", *Phys. Rev. B* **33**, 551-558, (1986).
4. U. Sivan and Y. Imry, "Energy Level Correlation Function and ac Conductivity of a Finite Disordered System", *Phys. Rev. B* **35**, 6074-6083 (1987).
5. U. Sivan and A. Seitz, "Lightwaves Localization in Dielectric Waveguides", *Europhys. Lett.* **5**, 139-144 (1988).
6. U. Sivan, O. Entin-Wohlman, and Y. Imry, "Orbital Magnetoconductance in the Variable Range Hopping Regime", *Phys. Rev. Lett.* **60**, 1566-1569 (1988).
7. U. Sivan, R. Blumenthal, Y. Meir, and O. Entin-Wohlman, "Dynamic Structure Factor of a Deterministic Fractal", *Europhys. Lett.* **7**, 249-253 (1988).

8. U. Sivan and Y. Imry, "de Haas-van Alphen and Aharonov-Bohm-type Persistent Current Oscillations in Singly Connected Quantum Dots", *Phys. Rev. Lett.* **61**, 1001-1004 (1988).
9. U. Sivan, Y. Imry, and C. Hartzstein, "Aharonov-Bohm and Quantum Hall Effects in Singly Connected Quantum Dots", *Phys. Rev. B* **39**, 1242-1250 (1989).
10. O. Entin-Wohlman, Y. Imry, and U. Sivan, "Orbital Magnetoconductance in the Variable Range Hopping Regime", *Phys. Rev. B* **40**, 8342-8348 (1989).
11. O. Entin - Wohlman, U. Sivan, R. Blumenfeld, and Y. Meir, "Dynamic Structure Factor of a Fractal", *Physica D* **38**, 93-97 (1989).
12. U. Sivan, M. Heiblum, and C. P. Umbach, "Hot Ballistic Transport and Phonon Emission in a Two-Dimensional Electron Gas", *Phys. Rev. Lett.* **63**, 992-995 (1989).
13. C. P. Umbach, A. Palevski, M. Heiblum, and U. Sivan, "Lateral Tunneling and Ballistic Transport in Two-Dimensional Electron Gas Devices Defined by Nanostructure Gates", *J. Vac. Sci. Technol. B* **7**, 2003 (1989).
14. U. Sivan, M. Heiblum, C. P. Umbach, and H. Shtrikman, "Electrostatic Electron Lens in the Ballistic Regime", *Phys. Rev. B* **41**, 7937-7940 (1990).
15. I. Kander, Y. Imry, and U. Sivan, "Effects of Channel Opening and Disorder on the Conductance of Two Dimensional Wires", *Phys. Rev. B* **41**, 12941-12944 (1990).
16. B. Laikhtman, U. Sivan, A. Yacoby, C. P. Umbach, M. Heiblum, J. A. Kash, and H. Shtrikman, "Long Mean Free Path of Hot Electrons Injected to Higher Subbands", *Phys. Rev. Lett.* **65**, 2181-2184 (1990).
17. D. Kowal, U. Sivan, O. Entin-Wohlman, and Y. Imry, "Transmission in Multiply Connected Wire Systems", *Phys. Rev. B* **42**, 9009-9018 (1990).
18. U. Sivan, A. Palevski, M. Heiblum, and C. P. Umbach, invited paper on "Ballistic Transport in a Two-Dimensional Electron Gas", *Solid State Electronics*, **33**, 979-986 (1990).
19. M. Heiblum, A. Palevski, U. Sivan, and C. P. Umbach, "Hot Electron Transport in the Plane", *Surf. Sci.* **229**, 155 (1990).
20. U. Sivan, O. Entin-Wohlman, and Y. Imry, "Orbital Magnetoconductance in the Variable Range Hopping Regime - Percolation Approach" pp. 151-167, in "Hopping and Related Phenomena", edited by Hellmut Fritzsche and Michael Pollak, 1990 World Scientific publishing company.
20. A. Palevski, U. Sivan, M. Heiblum, C. P. Umbach, and H. Shtrikman, "Hot Electron Transport in Two - Dimensional Structures", *Acta Physica Polonica A* **79**, no.1, 59-69 (1991).

21. A. Yacoby, U. Sivan, C. P. Umbach, and J. M. Hong, "Interference and Dephasing by Electron - Electron Interaction on Length Scales Shorter Than The Elastic Mean Free Path", *Phys. Rev. Lett.* **66**, 1938-1941 (1991).
22. U. Sivan, A. Yacoby, C. P. Umbach, and J. M. Hong, "Coherence and Electron - Electron Interaction in Ballistic Conductors", in "Nanostructures and Mesoscopic Systems", pp. 119-130, Academic Press Inc. (1992).
23. M. Heiblum and U. Sivan "Hot Electron Transport in a High Mobility Two Dimensional Electron Gas", in "Hot Carriers in Semiconductors Physics and Applications", edited by J. Shah, Academic Press Inc. (1992).
24. U. Sivan, P. M. Solomon, and H. Shtrikman, "Coupled Electron - Hole Transport", *Phys. Rev. Lett.* **68**, 1196-1199 (1992).
25. U. Sivan, K. Milkove, F. P. Milliken, S. Rishon, D. Kern, "Spectroscopy, Electron-Electron Interaction, and Energy Level Statistics in a Disordered Quantum Dot", *Physica Scripta*, **49B**, 446-448 (1993).
26. S. A. Rishon, Y. H. Lee, K. R. Milkove, J. M. Hong, V. Boegli, M. DeFranza, U. Sivan, and D. P. Kern, "Integrated Approach to Quantum Dot Fabrication", *J. Vac. Sci. Technol. B*, **11**, 2607-2611 (1993).
27. U. Sivan, K. Milkove, F. P. Milliken, S. Rishon, D. Kern, "Spectroscopy, Electron-Electron Interaction, and Energy Level Statistics in a Disordered Quantum Dot", *Europhysics Lett.* **23**, 605-611 (1994).
28. U. Sivan, Y. Imry, and A. G. Aronov, "Quasi Particle Lifetime in a Quantum Dot", *Europhys. Lett.* **28**, 115-120 (1994).
29. Y. Imry and U. Sivan, Invited review paper on "Recent Developments in Mesoscopic Physics", special issue on *Highlights in Condensed Matter Physics*, *Solid State Comm.*, **92**, 83-87 (1994).
30. S. Shapira, U. Sivan, P. M. Solomon, E. Buchstab, and M. Tischler, "The Thermodynamics of a Charged Fermion Layer at High r_s Values", *Phys. Rev. Lett.* **77**, 3181-3184 (1996).
31. U. Sivan, R. Berkovits, Y. Aloni, O. Prus, A. Auerbach, and G. Ben-Yoseph, "Mesoscopic Fluctuations in the Ground State Energy of Disordered Quantum Dots", *Phys. Rev. Lett.* **77**, 1123-1126 (1996).
32. O. Prus, A. Auerbach, U. Sivan, Y. Aloni, and R. Berkovits, "Even-Odd Correlations in Capacitance Fluctuations of Quantum Dots", *Phys. Rev. B* **54**, R14289-92 (1996).
33. Y. Imry, A. Stern, and U. Sivan, "Electron-electron Scattering and Transport in Granular Systems", *Europhys. Lett.* **39**, 639-643 (1997).
34. R. Berkovits and U. Sivan, "Transmission Through an Interacting Quantum Dot", *Europhys. Lett.* **41**, 653-658 (1998).

35. E. Brann, Y. Eichen, U. Sivan and G. Ben Yoseph, "DNA templated assembly and electrode attachment of conducting silver wire", *Nature*, **391**, 775-778 (1998).
36. Y. Eichen, E. Brann, U. Sivan, and G. Ben Yoseph, "Self Assembly of Nanoelectronics Components and Circuits Using Biological Templates", *Acta Polymerica* **49**, 663-670 (1998).
37. Y. Yaish, Oleg Prus, Evgeny Buchstab, Shye Shapira, Gidi Ben Yoseph, Uri Sivan, and Ady Stern, "Interband Scattering and the "Metallic Phase" of Two Dimensional Holes in GaAs/AlGaAs", *Phys. Rev. Lett.* **84**, 4954-4957 (2000).
38. K. Keren, A. Stern, and U. Sivan, "Electron-Electron lifetime-What is the Difference Between an Atom and a Quantum Dot", *Eur. Phys. J. B* **18**, 311-318 (2000).
39. O. Prus, M. Reznikov, U. Sivan, and V. Pudalov, "On the Cooling of Electrons in a Silicon Inversion Layer", *Phys. Rev. Lett.* **88**(1) (2001).

Papers published in proceedings.

1. E. Kapon, U. Sivan, J. Salzman, and A. Katzir, "Heterodyne Detection Using Multimode Waveguide Y-Couplers", *SPIE Proceedings*, San Diego 1983.
2. U. Sivan, B. Lalkhtman, A. Yacoby, C. P. Umbach, M. Heiblum, J. A. Kesh, and H. Shtrikman, "Long Mean Free Path of Hot Electrons Selectively Injected to Higher Subbands", *Proceedings of the 20th International Conference on The Physics of Semiconductors*, E. M. Anastassakis and J. D. Joannopoulos editors, World Scientific (1990), pp. 375-378.
3. U. Sivan, P. M. Solomon, and H. Shtrikman, invited paper on "Coupled Electron - Hole Gases", *Proceedings of the 21st International Conference on the Physics of Semiconductors*, Beijing, China (1992).
4. P. M. Solomon, U. Sivan, and H. Shtrikman, "Transport in Coulombically coupled Electron-Hole System" (abstract), *proceedings of the 10th International Conference on the Physics of Two Dimensional Systems*, Newport, RI, USA (1993).
5. U. Sivan, invited paper on "Spectral Properties of Disordered Quantum Dots", *Proceedings of the XXIXth Rencontre the Moriond*, Villars-sur-Ollon, Switzerland (1994).
6. S. Shapira, U. Sivan, P. M. Solomon, E. Buchstab, M. Tischler, and G. Ben Yoseph, *Proceedings of 23rd International Conference on the Physics of Semiconductors*, World Scientific, Singapore, M. Schöffler and R. Zimmermann eds. P. 2311-2314 (1996).

-
7. S. Shapira, U. Sivan, P. M. Solomon, E. Buchstab, and M. Tischer, "The Thermodynamics of a Charged Fermion Layer at High r_s Values", in the proceedings of the 11th International Conference on the Electronic Properties of Two-Dimensional Systems, Nottingham, UK, Surf. Sci. (Netherlands), vol.361-362, p. 113-116 (1996).

F. PATENTS

Four patent applications on applications of interfacing molecular biology with molecular electronics.

DNA-templated assembly and electrode attachment of a conducting silver wire

Erez Braun*, Yoav Eichen†, Uri Sivan*‡ & Gdalyahu Ben-Yoseph*‡

* Department of Physics, † Department of Chemistry, ‡ Solid State Institute, Technion-Israel Institute of Technology, Haifa 32000, Israel

Recent research in the field of nanometre-scale electronics has focused on two fundamental issues: the operating principles of small-scale devices, and schemes that lead to their realization and eventual integration into useful circuits. Experimental studies on molecular¹ to submicrometre² quantum dots and on the electrical transport in carbon nanotubes^{3–5} have confirmed theoretical predictions^{6–8} of an increasing role for charging effects as the device size diminishes. Nevertheless, the construction of nanometre-scale circuits from such devices remains problematic, largely owing to the difficulties of achieving inter-element wiring and electrical interfacing to macroscopic electrodes. The use of molecular recognition processes and the self-assembly of molecules into supramolecular structures^{9,10} might help overcome these difficulties. In this context, DNA has the appropriate molecular-recognition¹¹ and mechanical^{12–16} properties, but poor electrical characteristics prevent its direct use in electrical circuits. Here we describe a two-step procedure that may allow the application of DNA to the construction of functional circuits. In our scheme, hybridization of the DNA molecule with surface-bound oligonucleotides is first used to stretch it between two gold electrodes; the DNA molecule is then used as a template for the vectorial growth of a 12 μm long, 100 nm wide conductive silver wire. The experiment confirms that the recognition capabilities of DNA can be exploited for the targeted attachment of functional wires.

The first step in the construction of the silver wire involves the self-assembly of a DNA template connecting two gold electrodes 12–16 μm apart (see Fig. 1 for an outline of the procedure). First, 12-base oligonucleotides, derivatized with a disulphide group at their 3' end, are attached to the electrodes through sulphur–gold interactions. The electrodes are each marked with specific but different oligonucleotide sequences. A connection is then made by hybridizing two distant surface-bound oligonucleotides with a 16 μm long and fluorescently labelled λ -DNA that contains two 12-base sticky ends, where each of the ends is complementary to one of the two different sequences attached to the gold electrodes. Hybridization on both ends is facilitated by covering the electrodes with an aqueous solution containing the λ -DNA and inducing a flow perpendicular to the electrodes, thereby stretching the λ -DNA molecules in the flow direction (other stretching methods can be used; for application of an electric field, see ref. 17). The flow is terminated when a single DNA bridge is observed by fluorescence microscopy (see Fig. 2), usually after a few minutes. Curving of the

DNA bridge under a flow parallel to the electrodes shows it to be attached solely to the electrodes. The method does not guarantee a single DNA bridge. However, *in situ* video microscopy and imaging of the resulting silver wire by atomic force microscopy (AFM; see below) reveal a silver bridge only in places where DNA was previously fluorescently imaged. We also tried stretching the DNA between two electrodes in the reverse order, performing hybridization and ligation of the disulphide-derivatized oligonucleotides to the long DNA molecule before it was applied to the sample (see Methods section). The binding of the derivatized λ -DNA in this case was again aided by an induced perpendicular flow. Both methods work equally well.

To instill electrical functionality, silver metal is vectorially deposited along the DNA molecule. The three-step chemical deposition process (see Fig. 1 and Methods) is based on selective localization of silver ions along the DNA through Ag^+/Na^+ ion-exchange¹⁸ and formation of complexes between the silver and the DNA bases^{19–21}. The Ag^+/Na^+ ion-exchange process is monitored by following the almost instantaneous quenching of the fluorescence signal of the labelled DNA. The ion-exchange process, which is highly selective and restricted to the DNA template only, is terminated when the fluorescence signal drops to 1–5% of its initial value (the quenching is much faster than normal bleaching of the fluorescent dye). The silver ion-exchanged DNA is then reduced to form nanometre-sized metallic silver aggregates bound to the DNA skeleton. These silver aggregates are subsequently further 'developed', much as in the standard photographic procedure, using an acidic solution of hydroquinone and silver ions under low light conditions^{22,23}. Such




Figure 1 Construction of a silver wire connecting two gold electrodes. The top left image shows the electrode pattern (0.5 × 0.5 mm) used in the experiments. The two 50 μm long, parallel electrodes are connected to four (100 × 100 μm) bonding pads. **a**, Oligonucleotides with two different sequences attached to the electrodes. **b**, λ -DNA bridge connecting the two electrodes. **c**, Silver ion-loaded DNA bridge. **d**, Metallic silver aggregates bound to the DNA skeleton. **e**, Fully developed silver wire. A full description of the preparation steps can be found in the Methods section.

letters to nature

solutions are metastable, and spontaneous metal deposition is normally very slow. However, the silver aggregates on the DNA act as catalysts and significantly accelerate the process. Under the experimental conditions, metal deposition therefore occurs only along the DNA skeleton, leaving the passivated glass practically clean of silver. The silver deposition process is monitored *in situ* by differential interference contrast (DIC) microscopy and terminated when a trace of the metal wire is clearly observable under the microscope. The metal wire follows precisely the previous fluorescence image of the DNA skeleton. The structure, size and conduction properties of the metal wire are reproducible and dictated by the 'developing' conditions.

AFM images of segments of a 100 nm wide, 12 μm long wire connecting the two gold electrodes are presented in Fig. 3. The images are representative for the whole wire. As clearly seen, the wire consists of grains of 30–50 nm diameter deposited along the DNA skeleton. We have also fabricated wires as narrow as 25 nm but AFM imaging showed the silver coating to be discontinuous, with some gaps between silver grains.

Two-terminal electrical measurements were made on the silver wire shown in Fig. 3. The resultant I - V curves, obtained with an apparatus having an internal resistance of $\geq 10^{13} \Omega$, are displayed in Fig. 4. The curves are highly nonlinear and asymmetric with respect to zero bias. The shape of the curves depends on the voltage scan direction indicated by arrows. Approaching zero voltage from large positive or negative bias, a zero-current plateau develops with differential resistance larger than $10^{13} \Omega$. At a higher bias, the wire again becomes conductive with a differential resistance somewhat lower than in the original bias polarity. Repeated measurements of a given wire, in the same scan direction, yield reproducible I - V curves.

The origin of the extremely high resistance at small bias, and of the dependence of the current on the voltage scan direction, are not yet clear. If one invokes the Coulomb blockade phenomenon and the grain charging energy, ~ 0.1 eV, as inferred from the grain size in the AFM images, the large plateau requires simultaneous charging of a large number of grains in series. It is not clear, however, whether such a mechanism can yield a history-dependent I - V curve. Another source of nonlinearity might be inter-grain boundary resistance due to silver corrosion. The shape of the I - V curve could then be attributed to electrochemical processes in the corrosion barriers.

The length of the zero bias plateau in different wires may vary

from a fraction of a volt to roughly 10 V, depending on the silver growth conditions. The solid line in Fig. 4b depicts, for example, the I - V curve of a different wire in which the silver growth on the DNA was more extensive. Compared with the first wire, the plateau was reduced from about 10 V to 0.5 V, and the differential resistance at voltages beyond the plateau was reduced from 30 M Ω to 7 M Ω . By applying voltages higher than about 50 V to the wires that went through extensive silver deposition, the plateau could usually be permanently eliminated to give ohmic behaviour (dashed-dotted line in Fig. 4b).

Each experiment was accompanied by two control measurements to investigate whether either the DNA bridge or the deposited silver on their own could contribute to the observed electrical transport. The two insets in Fig. 4b show the I - V curves obtained for a λ -DNA bridge in the absence of silver deposition (bottom inset) and for a neighbouring device without a DNA bridge that underwent the full silver deposition treatment (top inset). Both control measurements consistently yielded a resistance higher than our measurement capabilities, $10^{13} \Omega$. (Note that the current scale in both insets is 100 times smaller than in the main graph.) Our direct d.c. measurements showed the 16 μm long double-stranded DNA to be practically insulating. It is not clear how to compare our measurements with previously published electron-transfer rates obtained from optical measurements on short DNA segments^{24–26}. Perfect correlation was found between the appearance of a DNA bridge in the fluorescence image, the formation of a silver wire connecting the




Figure 2 Fluorescence image of the DNA bridge. Fluorescently labelled (YoYo-1, Molecular Probes, Eugene, Oregon) λ -DNA is stretched between two gold electrodes (dark strips), 16 μm apart. The electrodes are connected to large bonding pads 0.25 mm away.




Figure 3 Atomic force microscopy images (Dimension 3000, Digital Instruments) of a silver wire connecting two gold electrodes 12 μm apart. **a**, 1.5 μm ; **b**, 0.5 μm field sizes. Note the granular morphology of the conductive wire.

electrodes, and the resulting conduction between the electrodes. These observations, in conjunction with the control experiments, prove that the electric current is carried solely by the silver deposited on the DNA bridge.

The remarkable recognition capabilities of DNA, used in this study to construct a metal wire connecting two electrodes, have also been exploited in other recent studies. The use of DNA has, for example, allowed researchers to organize colloidal particles into macroscopic crystal-like aggregates^{27,28} and to dictate the shape and size of semiconductor nanoparticle assemblies²⁹. The use of a DNA polyanion as a template for the assembly of electronic materials is not limited to metals. In another study, we have used a DNA template to fabricate a poly-(*p*-phenylene vinylene) (PPV) filament by attaching a positively charged *pre*-PPV polymer to the stretched DNA and subsequently treating it to form a highly photoluminescent PPV wire³⁰.

The construction of functional nanoscale electronic devices is likely to require self-assembly processes. The present study provides a step towards such developments: we have demonstrated that DNA is a sophisticated substrate for the targeted attachment of a conductive metal wire whose width is well below the minimal dimensions accessible by standard, large-scale microelectronics technology. Even though the hysteric *I*-*V* curves that we observe are interesting, low-resistance, ohmic metal wires are essential for

future applications. Such wires might be accessible by making use of different growth conditions, different metal types or post-growth thermal treatments. □

Methods

Sample preparation. A glass coverslip is immersed in fuming nitric acid for 10 min, rinsed with DI water, immersed in a 1M NaOH solution for a further 10 min and rinsed again with DI water. The cleaned glass is dried thoroughly and then passivated against spurious DNA binding by immersion for 12 hours in a 1:5 v/v solution of trimethyl chlorosilane (Sigma) in tetrachloroethane (Sigma). The sample is then rinsed carefully several times with tetrachloroethane and isopropanol and dried thoroughly. Electrodes, 12 μ m or 16 μ m apart, are defined on the coverslip by standard photolithography and subsequent high vacuum deposition of a 50 nm gold layer on top of a 10 nm titanium adhesion layer. A lift-off process then follows.

Constructing the DNA bridge (see Fig. 1). One gold electrode is wetted with a 10⁻⁴ μ l droplet of an aqueous solution containing 0.2 nM of a 5'-GGGCGGCGACCT-3'-disulphide oligonucleotide (Oligo A) and 10 mM NaCl. Similarly, the other electrode is marked with a solution containing 5'-AGGTCCGCGCCC-3'-disulphide oligonucleotides (Oligo B). Both oligonucleotides are synthesized using a 3'-C6-disulphide CPG (Clontech Laboratories, Palo Alto, California). After rinsing, the structure is covered by a 100 μ l solution of λ -DNA (0.2 pM, Promega, Madison, Wisconsin) in 10–100 mM NaCl. The λ -DNA has two sticky ends that are complementary to Oligos A and B. A flow perpendicular to the electrodes is induced by micropipette suction of the solution. The flow stretches the λ -DNA molecules perpendicular to the electrodes, leading to their hybridization with Oligos A and B attached to the two electrodes.

Hybridization and ligation before application to the electrodes. The two types of oligonucleotides are phosphorylated at their 5' ends using T4 polynucleotide kinase (New England Biolabs, Beverly, Massachusetts). The 5' phosphate of the λ -DNA is removed using Shrimp alkaline phosphatase (Amersham, Arlington Heights, Illinois) to prevent ligation of its complementary sticky ends to form a cyclic DNA. One type of oligonucleotide is then hybridized with the λ -DNA (10³-fold excess of oligonucleotides) while the solution (sodium phosphate buffer and 1M NaCl, pH = 7) slowly cools down (16 hours) from 75 °C to 4 °C. The hybridization is followed by ligation using T4 ligase (New England Biolabs) at 16 °C for 16 hours. The solution is then filtered (Microcon-100, Amicon, Beverly, Massachusetts) to remove excess free oligonucleotides. The second type of oligonucleotide is then hybridized and ligated with the other sticky end of the λ -DNA using the same procedures.

Silver deposition. Being a polyanion, the DNA bridge is loaded with silver ions by Na⁺/Ag⁺ ion exchange using a 0.1M AgNO₃ basic aqueous solution (ammonium hydroxide, pH = 10.5). The silver ion/DNA complex is reduced using a basic hydroquinone solution (0.05M, ammonium hydroxide, pH = 10.5) to form small metallic silver aggregates bound to the DNA skeleton. The DNA templated wire is 'developed' using an acidic solution (pH = 3.5, citrate buffer) of hydroquinone (0.05M) and silver ions (0.1M) under low light conditions to give a silver wire along the DNA skeleton.

Imaging setup. An inverted microscope (Axiovert 135, Zeiss), equipped with 100 \times oil-immersed objective, 100-W mercury lamp with filters, image intensifier (C-2400 Hamamatsu) and video processor (Omnex, Imagen Instrumentation), is used for fluorescence imaging of the DNA molecules. The same microscope and objective, with the proper DIC slide, and a 100-W halogen lamp are used for DIC imaging of the silver wire.

Electrical measurements. All electrical measurements were carried out between two bonding pads (see Fig. 1) using a 4145 HP parameter analyser with internal resistance $\geq 10^{10}$ Ω and current resolution of 10 fA.

Received 11 July; accepted 4 November 1997.

1. Porath, D. & Mülloh, O. Single electron tunnelling and level spectroscopy of isolated C₆₀ molecules. *J. Appl. Phys.* **81**, 1–4 (1997).
2. Mehner, U. & Foxman, E. B. Single-electron phenomena in semiconductors. *Semiconductor Sci. Technol.* **11**, 253–284 (1996).
3. Langer, L. et al. Quantum transport in a multiwalled carbon nanotube. *Phys. Rev. Lett.* **76**, 479–482 (1996).
4. DNA, S. J. et al. Individual single-wall carbon nanotubes as quantum wires. *Nature* **386**, 474–477 (1997).
5. Bockrath, M. et al. Single-electron transport in ropes of carbon nanotubes. *Science* **273**, 1922–1925 (1997).




Figure 4 Experimentally observed *I*-*V* curves. **a**, Two terminal *I*-*V* curves of the silver wire shown in Fig. 3. Arrows indicate voltage scan direction. The two curves in each direction present repeated measurements thus demonstrating the stability of a given wire. Note the different asymmetry pertaining to the two scan directions. **b**, *I*-*V* curves of a different silver wire in which the silver growth was more extensive than in **a**. Extensive growth resulted in a narrower current plateau (solid curve), on the order of 0.5 V, and a lower differential resistance (7 M Ω versus 30 M Ω in **a**). By applying 50 V to the wire, the plateau has been permanently eliminated to give an ohmic behaviour (dashed-dotted line), over the whole measurement range. *I*-*V* curves of a DNA bridge with no silver deposition, and silver deposition without a DNA bridge, are depicted in the bottom and top insets, respectively. In both cases, the sample is insulating.

letters to nature

6. Averin, D. V. & Likharev, K. K. In *Atomic Scale Phenomena in Solids* (eds Altshuler, B., Lee, R. & Webb, R.) 173 (Elsevier, Amsterdam, 1991).
7. Grabert, M. & Devoret, M. (eds) *Single Charge Tunneling* (Plenum, New York, 1992).
8. Kastner, M. A. The single-electron transistor. *Rev. Mod. Phys.* **64**, 849–858 (1992).
9. Lehn, J. M. *Supramolecular Chemistry: Concepts and Perspectives* (VCH, Weinheim, 1995).
10. Atwood, J. L. et al. (eds) *Comprehensive Supramolecular Chemistry* (Pergamon, Oxford, 1996).
11. Watson, J. D. et al. *Molecular Biology of the Gene* 4th edn (Benjamin Cummings, Menlo Park, 1987).
12. Austin, R. H. et al. Stretching genes. *Phys. Today* **50**, 32–38 (1997).
13. Benishvili, D. et al. Stretching DNA with a recording semiconducting diode. *Phys. Rev. Lett.* **74**, 4734–4737 (1995).
14. Benishvili, D. et al. Alignment and sensitive detection of DNA by a moving interface. *Science* **265**, 2094–2096 (1994).
15. Wirtz, D. Direct measurement of the transport properties of a single DNA molecule. *Phys. Rev. Lett.* **75**, 2436–2439 (1995).
16. Zimmerman, R. M. & Cox, E. C. DNA stretching on functionalized gold surfaces. *Nucleic Acids Res.* **22**, 492–497 (1994).
17. Schwartz, D. C. & Kowal, M. Conformational dynamics of individual DNA molecules during gel electrophoresis. *Nature* **338**, 520–522 (1987).
18. Barton, J. K. In *Inorganic Chemistry* (eds Bertini, I. et al.) ch. 8 (University Science Books, Mill Valley, 1994).
19. Spiro, T. G. (ed.) *Nucleic Acid–Metal Ion Interactions* (Wiley Interscience, New York, 1980).
20. Marchetti, L. G., Kistnermacher, T. J. & Rossi, M. *J. Am. Chem. Soc.* **99**, 2797–2798 (1977).
21. Eichorn, G. L. (ed.) *Inorganic Biochemistry* Vol. 2, ch. 33–34 (Elsevier, Amsterdam, 1973).
22. Holgate, C. S. et al. Immunogold-silver staining: new method of immunostaining with enhanced sensitivity. *Histochem. J.* **31**, 938–944 (1983).
23. Birrell, G. B. et al. Silver-enhanced colloidal gold as a cell surface marker for photoelectron microscopy. *J. Histochem. Cytochem.* **34**, 339–343 (1986).
24. Hall, S. D., Holmlund, B. E. & Barton, J. K. Oxidative DNA damage through long-range electron transfer. *Nature* **382**, 731–733 (1996).
25. Arkin, M. R. et al. Rates of DNA-mediated electron transfer between metallointercalators. *Science* **273**, 473–480 (1996).
26. Lenda, R. D. et al. Distance-dependent electron transfer in DNA hairpins. *Science* **277**, 673–676 (1997).
27. Mirkha, C. A. et al. A DNA-based method for rationally assembling nanoparticles into nanoscopic materials. *Nature* **382**, 607–609 (1996).
28. Alivisatos, A. P. et al. Organization of 'nanocrystal molecules' using DNA. *Nature* **382**, 609–611 (1996).
29. Coffey, J. L. et al. Dictation of the shape of molecular semiconducting nanoparticle assemblies by plasmid DNA. *Appl. Phys. Lett.* **67**, 3451–3453 (1996).
30. Burroughes, J. H. et al. Light-emitting diodes based on conjugated polymers. *Nature* **347**, 539–541 (1990).

Acknowledgements. We thank T. Naras, A. Adinolfi, W. Koplan and S. Lipson for discussions and technical assistance.

Correspondence should be addressed to E.B. (rc@physics.technion.ac.il).